

Prepared for *Theor. Chem. Acc.* (Fernando Bernardi memorial issue)

November 6, 2006

Improved direct diabatization and coupled potential energy surfaces for the photodissociation of ammonia

Zhen Hua Li, Rosendo Valero, and Donald G. Truhlar

Department of Chemistry and Supercomputing Institute,

University of Minnesota, Minneapolis, MN 55455-0431

Abstract. Diabatic potential energy surfaces are a convenient starting point for dynamics calculations of photochemical processes, and they can be calculated by the fourfold way direct diabatization scheme. Here we present an improved definition of the reference orbital for applying the fourfold way direct diabatization scheme to ammonia. The improved reference orbital is a geometry-dependent hybrid orbital that allows one to define consistent dominant configuration lists at all geometries important for photodissociation. Using diabatic energies calculated with the new reference orbital and consistent dominant configuration lists, we have refitted the analytical representations of the ground and the first electronically excited singlet-state potential energy surfaces and the diabatic coupling surface. Improved functional forms were used to reproduce the experimental dissociation energies and excitation energies, which will be important for subsequent simulations of photochemical dynamics. We find that the lowest-energy conical intersection point is at 5.16 eV, with C_{2v} symmetry.

1. Introduction

Nangia and one of the authors¹ have recently used the fourfold way direct diabaticization scheme²⁻⁴ to generate coupled potential energy surfaces (PESs) for the photodissociation of ammonia. This involved using the threefold density criterion and a reference orbital on nitrogen to generate diabatic molecular orbitals²⁻⁴ (DMOs), specification of diabatic prototypes in terms of lists of dominant configuration state functions (CSFs) defined in terms of the DMOs,²⁻⁴ and transformation of adiabatic many-electron state functions to diabatic state functions by the principle of configurational uniformity.²⁻⁶ The diabatic surfaces and couplings were calculated at 3600 geometries, fitted to economical analytic functions incorporating the full permutational symmetry of the three hydrogen atoms, and used to explore the four-dimensional conical intersection seams.

In subsequent unpublished work we attempted to employ this analytic potential energy matrix for semiclassical trajectory calculations, and this attempt showed that the surfaces had unrealistic values in certain regions of configuration space that were apparently not well enough represented in our data set (regions with partly dissociated bonds where the sets of bond angles were in certain ranges and also in certain high-energy regions where the ammonia umbrella is very closed). In the process of generating more data for a wider range of geometries, we discovered that we needed to develop a more robust definition of the reference orbital and a better fitting function to achieve a smooth global potential energy matrix with more accurate asymptotic behavior. Along the way we also developed improved dominant configuration lists for the diabatic states. We used the improved reference orbital and dominant configuration lists to generate diabatic surfaces and their couplings at about 8200 geometries spanning a broader range than in the previous work, and we fitted these surfaces to the new functional forms. We then tested the resulting potentials in sample semiclassical trajectory calculations to ensure that the nonphysical energies were properly corrected.

In the present article we report these new, more robust procedures and the resulting potential energy surfaces and couplings. A dynamical study employing these new surfaces is in progress.

2. Theory and Calculations

We will repeat as little as possible of the theory^{2–4} and the details of the previous¹ application to NH_3 . The MC-QDPT method^{7–9} with a state-averaged CASSCF reference function^{10,11} was used in all the calculations that were used for diabaticization. The $1s$ core orbital of nitrogen was kept frozen (that is, kept doubly occupied in all CSFs) in the MC-QDPT step, which includes all the double excitations from the active orbitals through second order. To avoid artifacts due to intruder states in the MC-QDPT wave function, the intruder state avoidance (ISA) method¹² was used in all the calculations. The level shift parameter¹² b of the ISA method was set to $0.02 E_h^2$ (note: $1 E_h \equiv 1$ hartree). The CASSCF wave function was obtained as a state average with equal weights for the lowest two adiabatic singlet electronic states of NH_3 . The active space contains eight electrons in seven orbitals, which is sometimes denoted (8,7). We used the 6-311++G(3df,3pd)^{13,14} basis set with six sets of six Cartesian d functions (three on N, one on each H) and one set of ten Cartesian f functions (on N).

The seven active orbitals were chosen as follows. First, we include in the active space the four occupied molecular orbitals (MOs) of the reference restricted Hartree-Fock (RHF) wave function, which are three bonding N–H MOs and one nonbonding orbital centered on the nitrogen atom. Then we include three virtual orbitals of the RHF wave function, in particular two antibonding N–H MOs and one Rydberg-type MO with $3s$ character centered on the nitrogen atom at short N–H distances. The Rydberg MO transforms into a $1s$ atomic orbital centered on the dissociating hydrogen atom at long N–H distances.

The electronic structure calculations used to generate the PESs were carried out with HONDOPLUS v.4.7.¹⁵ The ISA algorithm was added to HONDOPLUS from the version in GAMESS.¹⁶

Additional state-specific CASSCF calculations with the same active space and basis set were carried out for the minima of the ground and excited state PESs of NH_3 as well as for the $\text{NH}_2 + \text{H}$ dissociation limit with the NH_2 fragment in its ground and first excited electronic states. These calculations employed *Gaussian 03*.¹⁷

In order to establish the list of CSFs defining the diabatic prototypes of the fourfold way, first the diabatic MOs (DMOs) obtained by the threefold way at planar and nonplanar geometries and at short and long N–A distances (the dissociating hydrogen atom is denoted as H_A or A, depending on the context, and the other hydrogens are H_B and H_C or B and C) were correlated. It is only necessary to consider dissociation of one of the hydrogen atoms, since the two other possible N–H dissociations can be obtained by symmetry. The correlations are shown in Table 1. In this table, the NH_3 moiety is assumed to be oriented with the nitrogen atom at the origin and the two hydrogen atoms that do not dissociate situated in the xy plane. For planar geometries the dissociating hydrogen atom is also situated in the xy plane. A particular set of nonplanar geometries is defined such that the dissociating hydrogen atom is directed along the z axis. Although for a general nonplanar geometry the dissociating hydrogen atom can point in an arbitrary direction, this particular orientation of the dissociating hydrogen atom has been chosen to illustrate the problems encountered when one tries to define diabatic prototypes consistently for both planar and nonplanar geometries by applying the overlap criterion to the DMOs obtained by the threefold way.

In Table 1, for small N–A distances, “ $\sigma_{\text{N–A}}$ ” denotes a bonding DMO between the nitrogen atom and the dissociating hydrogen atom, “ p_z ” is a nonbonding p -type orbital centered on the nitrogen atom, “ R ” is a Rydberg-type $3s$ orbital centered on the nitrogen atom, and “ n ” is a nonbonding orbital centered on the nitrogen atom, but not oriented in a particular direction for nonplanar geometries. Likewise, for large N–A distances, “ sp^2 ” is a hybrid nonbonding DMO centered on the nitrogen atom and situated in the NH_2 fragment molecular plane, “ p_z ” is a nonbonding p -type orbital centered on the nitrogen atom, and “ h ” denotes the $1s$ orbital of the dissociated hydrogen atom.

In Table 1, note the interchange in the role of the first two DMOs between planar and some nonplanar geometries (with β less than about 75°) for intermediate N–A bond lengths (around 1.5–2.0 Å): for planar geometries, the $\sigma_{\text{N–A}}$ DMO is in the plane of the NH_2 moiety, whereas the nonbonding p_z DMO is perpendicular to that plane. However, for the abovementioned subset of nonplanar geometries, the $\sigma_{\text{N–A}}$ DMO is along the z axis and perpendicular to the NBC plane, with the nonbonding n DMO being now located in the NBC plane. The situation for nonplanar geometries is illustrated in Figure 1, where the n DMO is taken as the p_z orbital for the sake of illustration. Thus, the threefold way along with the overlap criterion for the DMOs at a sequence of geometries along a dissociation path would not provide a consistent list of CSFs for the diabatic prototypes at both planar and nonplanar geometries. In order to define consistent diabatic prototypes, it is necessary to use the more general fourfold way procedure. One reference nonbonding DMO was introduced, whose character transforms continuously between p_z , being perpendicular to the plane of the NBC moiety for planar geometries, and a nonbonding n orbital perpendicular to the ABC plane for nonplanar geometries. The new correlation of the DMOs is as shown in Table 2, where it can be seen that now the DMOs keep a uniform character at all geometries.

The need to define a reference orbital for the fourfold way procedure means that one has to choose a standard orientation for the NH_3 molecule. In the standard orientation, the nitrogen atom was placed at the origin with one hydrogen atom pointing along the positive x axis; the second hydrogen atom was situated in the xy plane with a positive value of y , and with the third hydrogen atom having an arbitrary orientation. A geometry-dependent reference DMO was defined as a DMO centered on the nitrogen atom and having nonbonding character at all molecular configurations. Thus, for separate $\text{NH}_2 + \text{H}$ fragments, the reference DMO is a p_z orbital centered on the nitrogen atom with coefficients computed by the threefold way at the following geometry of the NH_2 fragment: $r_2 = 1.5 \text{ \AA}$, $r_3 = 1.5 \text{ \AA}$, $\theta_1 = 120.0^\circ$, where r_2 is the distance from N to B, r_3 is the distance from N to C, and θ_1 is the BNC bond angle. We label the reference DMO at the dissociation limit as $\chi_{\text{ref},a}$. When all three N–H distances are short, the

reference DMO must point in the direction perpendicular to the plane formed by the three hydrogen atoms so as to keep its nonbonding character, even when the molecule does not have a plane of symmetry. In order to generate a reference DMO for an arbitrary geometry, two unit vectors, $\mathbf{v}^{(1)}$ and $\mathbf{v}^{(2)}$, were defined, where $\mathbf{v}^{(1)}$ points along the positive z direction and $\mathbf{v}^{(2)}$ points along the direction normal to the plane of the three hydrogen atoms. The components of these vectors are as follows:

$$v_x^{(1)} = 0, v_y^{(1)} = 0, v_z^{(1)} = 1 \quad (1)$$

$$v_x^{(2)} = \frac{y_{A-B} z_{B-C} - z_{A-B} y_{B-C}}{R_{A-B} R_{B-C} \sin \theta} \quad (2a)$$

$$v_y^{(2)} = -\frac{x_{A-B} z_{B-C} - z_{A-B} x_{B-C}}{R_{A-B} R_{B-C} \sin \theta} \quad (2b)$$

$$v_z^{(2)} = \frac{x_{A-B} y_{B-C} - y_{A-B} x_{B-C}}{R_{A-B} R_{B-C} \sin \theta} \quad (2c)$$

where \mathbf{R}_{A-B} and \mathbf{R}_{B-C} are bond vectors between atoms A and B and atoms B and C, respectively, and θ is the angle between these two vectors, x_{A-B} , y_{A-B} , and z_{A-B} are components of \mathbf{R}_{A-B} , and R_{A-B} is the magnitude of \mathbf{R}_{A-B} .

In order to guarantee a smooth, continuous transformation of the reference DMO from the strong-interaction region to separated product fragments, a switching function was defined as

$$F = \exp\left(-\left(r_1 - r_1^0\right)^2 / \Delta^2\right), \quad (3)$$

where r_1 is the distance from N to A, and r_1^0 and Δ are parameters discussed in the next paragraph. This switching function is then used to define a third vector, $\mathbf{v}^{(3)}$, as a linear combination of $\mathbf{v}^{(1)}$ and $\mathbf{v}^{(2)}$:

$$\mathbf{v}^{(3)} = F \mathbf{v}^{(2)} + (1 - F) \mathbf{v}^{(1)} \quad (4)$$

In Eq. (3), r_1^0 is a reference r_1 distance, and Δ is a parameter controlling the damping of the Gaussian function with distance. After some preliminary test calculations, the values $r_1^0 = 1.2 \text{ \AA}$ and $\Delta = 2.0 \text{ \AA}$ were found to give well-ordered DMOs along paths connecting planar with nonplanar geometries.

The final step is to apply $\mathbf{v}^{(3)}$ to the coefficients of the $\chi_{\text{ref},a}$ DMO in the contracted Gaussian basis set to obtain the required geometry-dependent reference DMO for the fourfold way. First one can define the triads of numbers $abc = 100, 010$ and 001 . Let $\phi_{\text{abc}}^{\text{CGTO}(1)}$, $\phi_{\text{abc}}^{\text{CGTO}(2)}$ and $\phi_{\text{abc}}^{\text{CGTO}(3)}$ denote the p -type normalized Cartesian contracted Gaussian-type orbitals (CGTOs) centered on the nitrogen atom in the 6-311++G(3df, 3pd) basis set.

$$\phi_{\text{abc}}^{\text{CGTO}(1)} = \sum_{i=1}^3 c_i x^a y^b z^c e^{-\zeta_i r^2} \quad (5a)$$

$$\phi_{\text{abc}}^{\text{CGTO}(2)} = x^a y^b z^c e^{-\zeta_4 r^2} \quad (5b)$$

$$\phi_{\text{abc}}^{\text{CGTO}(3)} = x^a y^b z^c e^{-\zeta_5 r^2} \quad (5c)$$

Likewise, let $\chi_{\text{ref},abc}^{\text{CGTO}(1)}$, $\chi_{\text{ref},abc}^{\text{CGTO}(2)}$ and $\chi_{\text{ref},abc}^{\text{CGTO}(3)}$ denote the coefficients of the reference DMO in the basis of these CGTOs. Finally, $v_{\text{abc}}^{(3)}$ are the components of the $\mathbf{v}^{(3)}$ vector. The equations defining the reference DMO are:

$$\chi_{\text{ref},abc}^{\text{CGTO}(1)} = v_{\text{abc}}^{(3)} \chi_{\text{ref},a,001}^{\text{CGTO}(1)} \quad (5d)$$

$$\chi_{\text{ref},abc}^{\text{CGTO}(2)} = v_{\text{abc}}^{(3)} \chi_{\text{ref},a,001}^{\text{CGTO}(2)} \quad (5e)$$

$$\chi_{\text{ref},abc}^{\text{CGTO}(3)} = v_{\text{abc}}^{(3)} \chi_{\text{ref},a,001}^{\text{CGTO}(3)} \quad (5f)$$

Recall from the definition of $\chi_{\text{ref},a}$ that it has nonzero coefficients only for the p_z -type CGTOs.

The diabatic molecular orbitals (DMOs) obtained by the fourfold way have the following character: $u_1 \equiv \sigma_{\text{N-B}}$, $u_2 \equiv \sigma_{\text{N-C}}$, $u_3 \equiv n/p_z$, $u_4 \equiv \sigma_{\text{N-C}/sp^2}$, $u_5 \equiv R/h$, $u_6 \equiv \sigma_{\text{N-B}}^*$, $u_7 \equiv \sigma_{\text{N-C}}^*$.

The notation $\text{DMO}_1 / \text{DMO}_2$ refers to the character of that particular DMO at short and long N–A bond lengths, respectively. In terms of these DMOs, the dominant CSFs defining each of the three diabatic prototypes, χ_1, χ_2, χ_3 for the ground state and χ_4, χ_5, χ_6 for the excited state, are:

$$\begin{aligned}
 \chi_1: & (u_1)^2 (u_2)^2 (u_3)^2 (u_4)^2 (u_5)^0 (u_6)^0 (u_7)^0, \\
 \chi_2: & (u_1)^2 (u_2)^2 (u_3)^2 (u_4)^1 (u_5)^1 (u_6)^0 (u_7)^0, \\
 \chi_3: & (u_1)^2 (u_2)^2 (u_3)^2 (u_4)^0 (u_5)^2 (u_6)^0 (u_7)^0, \\
 \chi_4: & (u_1)^2 (u_2)^2 (u_3)^1 (u_4)^2 (u_5)^1 (u_6)^0 (u_7)^0, \\
 \chi_5: & (u_1)^2 (u_2)^2 (u_3)^1 (u_4)^1 (u_5)^2 (u_6)^0 (u_7)^0, \\
 \chi_6: & (u_1)^2 (u_2)^1 (u_3)^2 (u_4)^2 (u_5)^1 (u_6)^0 (u_7)^0.
 \end{aligned} \tag{6}$$

The fourfold way procedure provides a smooth diabatic matrix in most of the dynamically relevant configuration space. The exceptions are some geometries where at least two of the N–H distances are relatively long (equal to or longer than 1.5 Å), when the trisector angle β is smaller than about 60° and for certain values of the HNH angles. Those points were removed from the global fit.

The coordinates used for the grids and the fitting are defined in Fig. 2. Note that in addition to the bond angles $\theta_1, \theta_2,$ and θ_3 , Fig. 2 shows the projected bond angles θ'_1, θ'_2 and θ'_3 that were originally introduced by Léonard et al.¹⁸ In addition we use the trisector angle β also introduced by Léonard et al.¹⁸ Most of the points for the fitting were contained in a regular grid constructed in coordinates $(r_1, r_2, r_3, \theta'_1, \theta'_2, \theta'_3, \beta)$ defined in Figure 2 as follows: r_1 was varied between 0.8 and 5.0 Å, and r_2 and r_3 were varied between 0.8 and 1.5 Å for a total of 44 different (r_1, r_2, r_3) values, which were combined with 182 sets of the four angles $(\theta'_1, \theta'_2, \theta'_3, \beta)$. These sets of four angles were obtained by combining 14 sets of projected bond angles $(\theta'_1, \theta'_2, \theta'_3)$ with 13 values of the trisector angle β , which is varied between 30° and 90° inclusive with a 5-degree regular increment. The groups of $(\theta'_1, \theta'_2, \theta'_3)$ angles are the following: (120,120,120); (120,150,90); (180,120,60); (120,90,150); (90,150,120); (180,90,90);

(150,90,60); (90,120,150); (180,60,120); (150,60,90); (120,60,60); (150,60,150); (150,90,120); and (150,120,90). Thus, the grid contains a total of 8008 points. The ranges of the coordinates and diabatic energies are shown in Table 3. The diabatic potential energies and couplings were plotted versus the trisector angle β and those points showing nonsmooth behavior were removed from the fit. All of the removed points had β angles less than 60° , as explained above. Some duplicated points, that is, points with different values of the coordinates defined above but with the same set of internuclear distances were also removed from the fit. After these operations, the remaining 5624 points were included in the fit.

The diabatic potential energy matrix was computed at additional sets of points to improve the fitting in particular regions of configuration space. A total of 2812 additional points with the coordinates and diabatic energies presented in rows 2–14 of Table 3 were computed, from which 2603 were included in the final fitting. The following numbers of points refer to those originally computed. Thus, 800 points were generated close to the ground-state NH_3 minimum (GS min in Table 3), with relatively small displacements from the equilibrium geometry, and 1200 additional points were chosen to map the region around the transition state for N–A dissociation on the V_2 PES (ES N–A saddle point in Table 3). In the labels, GS and ES denote ground state and excited state, respectively. To aid in the refinement of the frequencies of the NH_3 minimum on the V_2 PES, a normal mode analysis was carried out at the state-specific CASSCF(8,7)/6-311++G(3df, 3pd) level, and an additional 47 points were chosen along the six Cartesian normal mode vectors (ES min I and II in Table 3). Likewise, to assist in obtaining accurate stationary-point geometries and frequencies for the NH_2 fragments, 73 points suitable for the ground state and 48 points suitable for the excited state were added (NH_2 (\tilde{X}^2B_1) + H and NH_2 (\tilde{A}^2A_1) + H in Table 3).

Additional points were also chosen in repulsive regions of the PESs (Repulsive I–V in Table 3) to guide the fitting of the repulsive interactions (see Eq. (16) below). The sets of points I and II (about 200 points in total) were generated randomly in regions where preliminary fits generated spurious negative energies. The set of points III contains 20 points along the N–A

stretch curve with r_1 in the range 0.2–5.0 Å, and set IV contains 18 points with fixed N–H distances with θ_3' varying in such a way that one of the H–H distances decreases from 1.55 Å to about 0.25 Å. Finally, set V contains 100 points with very high energies (> 300 eV). An additional 312 points were computed to refine the long-distance tail of the PESs.

A total of 8227 points were included in the fitting.

Since the two diabatic surfaces cross at certain geometries, we will name the ground adiabatic potential energy surface as V_1 , the first singlet-excited adiabatic potential surface as V_2 , the diabatic potential energy surface with lower energy around the ground-state equilibrium geometry of NH₃ as U_{11} , and the higher-energy one as the U_{22} surface. Throughout the entire article, the zero of energy is at the equilibrium geometry of the ground adiabatic state of NH₃. Thus all values of the adiabatic surfaces V_1 and V_2 and the diabatic U_{11} and U_{22} are positive except $V_1 = 0$ at the lowest energy geometry.

3. Stationary-Point Data

Theoretical state-specific CASSCF as well as experimental (or best-estimate theoretical) data^{19–30} are presented in Tables 4 and 5. The data in the tables serve two purposes, namely (1) they are used to constrain the fitted PESs to reproduce the experimental NH₃ (\tilde{X}^1A_1) \rightarrow NH₂ (\tilde{X}^2B_1) + H and NH₃ (\tilde{X}^1A_1) \rightarrow NH₂ (\tilde{A}^2A_1) + H dissociation energies and the NH₃ (\tilde{X}^1A_1) \rightarrow NH₃ (\tilde{A}^1A_2'') and NH₂ (\tilde{X}^2B_1) \rightarrow NH₂ (\tilde{A}^2A_1) excitation energies (see below), and (2) they are used to provide a comparison of the *ab initio* and fitted geometry and frequencies of the NH₃ minima and of the NH₂ fragment in their electronic ground and first excited states.

The dissociation energies (D_0) and the excitation energies (T_0) at 0 K have been directly measured with good accuracy. To obtain the equilibrium dissociation energies D_e and the equilibrium excitation energies T_e (i.e., the energy differences on the PESs excluding the zero-point vibrational energies (ZPEs)), one needs to estimate anharmonic ZPEs.

The D_0 of the reaction NH₃ (\tilde{X}^1A_1) \rightarrow NH₂ (\tilde{X}^2B_1) + H has been extensively studied.³¹ Here we adopt the lower experimental photodissociation value of 4.602 ± 0.002 eV

determined by Mordaunt et al.,²³ which claims the highest accuracy and agrees very well with the latest compilation of Ruscic et al.³¹ Mordaunt et al. also calculated the D_e of this reaction (5.023 ± 0.003 eV). Due to the lack of experimental harmonic vibrational frequencies and their anharmonic couplings for the NH_2 ($\tilde{X}^2\text{B}_1$) radical, they used the following equation to obtain the ZPE of the NH_2 ($\tilde{X}^2\text{B}_1$) radical:

$$ZPE(\text{NH}_2) = ZPE(\text{NH}_2)^{\text{Theor}} \frac{ZPE(\text{NH}_3)^{\text{Exp}}}{ZPE(\text{NH}_3)^{\text{Theor}}} \quad (7)$$

where the theoretical ZPEs are calculated with purely theoretical anharmonic parameters. However, the theoretical parameters were calculated at different levels of theory; in particular MP4/6-31G(d,p) was used for ammonia³² while multireference configuration-interaction was used for the NH_2 ($\tilde{X}^2\text{B}_1$) radical. Here we adopt the value of Handy et al.,²² 0.925 eV (7461.53 cm^{-1}) for the ZPE of NH_3 ($\tilde{X}^1\text{A}_1$); this agrees very well with the value, 0.925 eV (7460.93 cm^{-1}) of Gatti et al.³³ For the ZPE of NH_2 ($\tilde{X}^2\text{B}_1$), we adopt the high level theoretical value of 0.511 eV of Gabriel et al.,²⁷ which agrees very well with other high-level theoretical results, in particular, 0.512 eV of Martin et al.²¹ and 0.512 eV of Demaison et al.³⁴ With these new choices of data, the D_e of the reaction NH_3 ($\tilde{X}^1\text{A}_1$) \rightarrow NH_2 ($\tilde{X}^2\text{B}_1$) + H is then calculated to be 5.016 eV, which is very close to the 5.023 eV given by Mordaunt et al.²³ We will use the value 5.016 eV.

To calculate D_e of the reaction NH_3 ($\tilde{X}^1\text{A}_1$) \rightarrow NH_2 ($\tilde{A}^2\text{A}_1$) + H, one needs to obtain the T_e excitation energy for NH_2 ($\tilde{X}^2\text{B}_1$) \rightarrow NH_2 ($\tilde{A}^2\text{A}_1$). The T_0 excitation energy has been accurately determined as 1.379 eV (11122.23 ± 0.05 cm^{-1}).³⁰ Since the ZPE of NH_2 ($\tilde{A}^2\text{A}_1$) corrected for anharmonicity is not directly available from the literature due to the lack of harmonic force constants and their anharmonic couplings, we used a similar strategy as Mordaunt et al. to estimate it:

$$ZPE(\text{NH}_2, \tilde{A}^2\text{A}_1) = ZPE(\text{NH}_2, \tilde{X}^2\text{B}_1) \frac{\left(ZPE^{\text{F}} + ZPE^{\text{H}} \right)_{\text{NH}_2, \tilde{A}^2\text{A}_1}}{\left(ZPE^{\text{F}} + ZPE^{\text{H}} \right)_{\text{NH}_2, \tilde{X}^2\text{B}_1}} \quad (8)$$

where

$$ZPE^F = \frac{1}{2} \sum_i \nu_i, \quad (9)$$

$$ZPE^H = \frac{1}{2} \sum_i \omega_i, \quad (10)$$

where ν_i is a fundamental frequency, and ω_i is a harmonic frequency. The use of Eq. (8) is based on the fact that $\frac{1}{2}(ZPE^F + ZPE^H)$ is a good approximation to the accurate ZPE.³⁵ The ZPE of NH_2 ($\tilde{\text{A}}^2\text{A}_1$) is calculated to be 0.529 eV by this procedure. With this value, the NH_2 ($\tilde{\text{X}}^2\text{B}_1$) \rightarrow NH_2 ($\tilde{\text{A}}^2\text{A}_1$) T_e excitation energy is calculated to be 1.361 eV, and a D_e of 6.377 eV for the NH_3 ($\tilde{\text{X}}^1\text{A}_1$) \rightarrow NH_2 ($\tilde{\text{A}}^2\text{A}_1$) + H reaction is obtained.

The fourth important energy difference that we try to reproduce with the analytical function is the NH_3 ($\tilde{\text{X}}^1\text{A}_1$) \rightarrow NH_3 ($\tilde{\text{A}}^1\text{A}_2''$) T_e excitation energy. The T_0 of this excitation has been determined as 5.722 eV (46150 cm^{-1}) by Henck et al.²⁴ and 5.719 eV (46130 cm^{-1}) by Douglas.³⁶ Henck et al. also estimated T_e using the *ab initio* ZPE of NH_3 ($\tilde{\text{X}}^1\text{A}_1$) calculated by Hargiss and Ermler³² and the experimental ZPE of NH_3 ($\tilde{\text{A}}^1\text{A}_2''$) that they determined (0.724 eV, 5840 cm^{-1}), which gives a value of 5.960 eV (48070 \pm 200 cm^{-1}). Using the best available ZPE for NH_3 ($\tilde{\text{X}}^1\text{A}_1$), which is 0.925 eV (7461.53 cm^{-1})²² as determined by Handy et al., T_e is calculated to be 5.923 eV, slightly lower than the value given by Henck et al.,²⁴ but closer to the best available theoretical value of 5.902 eV calculated by Bach et al.²⁵ We will use the value of 5.923 eV.

4. Functional forms and fitting procedures

4.1. Coordinate system used

Ten internal coordinates are used to fit the two potential energy surfaces and their coupling. In particular we used three N–H bond distances (r_1, r_2, r_3), three HNH bond angles ($\theta_1, \theta_2, \theta_3$), three H–H bond distances (r_4, r_5, r_6), and a trisector angle β (Figure 2). The trisector angle was originally introduced in earlier studies in order to better describe the inversion potential in the ground electronic state.^{18,22} The three H–H bond distances are used to add a

repulsive interaction (see Sec 4.2) at very small H-N-H bond angles to correct spurious nonphysical energies of the previous fitting functions. Notice that we used $(r_1, r_2, r_3, \theta_1', \theta_2', \theta_3', \beta)$ for the regular grids, but we used $(r_1, r_2, r_3, r_4, r_5, r_6, \theta_1, \theta_2, \theta_3, \beta)$ for fitting.

4.2. Functional forms

The U_{11} surface was fitted to the following functional form:

$$E(\mathbf{Q}) = \sum_{\substack{i \leq j \leq k \\ l \leq m \leq n \\ p}} C_{ijklmnp}^{(1)} \sum_{\alpha} A_{ijklmnp}^{(1)}(\mathbf{P}_{\alpha} \mathbf{Q}) + F(r_1, r_2, r_3, r_4, r_5, r_6) + E_D(r_1, r_2, r_3, r_4, r_5, r_6) \quad (11)$$

where \mathbf{P}_{α} is the permutation operator used in the previous study,¹ α (which equals 1, 2, ..., 6) labels the permutations of hydrogen atoms A, B, and C, and

$$A_{ijklmnp}^{(1)}(\mathbf{Q}) = B_i(r_1)B_j(r_2)B_k(r_3)g(\theta_1, r_2, r_3)g(\theta_2, r_1, r_3)g(\theta_3, r_1, r_2)h(p, \beta, r_1, r_2, r_3) \quad (12)$$

$$B_i(r) = \left[1 - e^{-\gamma^{(1)}(r - r_0^{(1)})} \right]^i \quad (13)$$

$$g(\theta_1, r_2, r_3) = \left\{ [\cos(\theta_0) - \cos(\theta_1)] e^{-\delta^{(1)}(r_2^2 + r_3^2)} \right\}^l \quad (14a)$$

$$g(\theta_2, r_1, r_3) = \left\{ [\cos(\theta_0) - \cos(\theta_2)] e^{-\delta^{(1)}(r_1^2 + r_3^2)} \right\}^m \quad (14b)$$

$$g(\theta_3, r_1, r_2) = \left\{ [\cos(\theta_0) - \cos(\theta_3)] e^{-\delta^{(1)}(r_1^2 + r_2^2)} \right\}^n \quad (14c)$$

$$h(p, \beta, r_1, r_2, r_3) = 1 + \left\{ 1 - \cos \left[p \left(\beta - \frac{1}{2} \right) \right] \right\} e^{-\delta^{(1)}(r_1^2 + r_2^2 + r_3^2)} \quad (15)$$

$$F(r_1, r_2, r_3, r_4, r_5, r_6) = B_{\text{NH1}}^{(1)} \sum_{i=1}^3 \frac{e^{-\beta_{\text{NH1}}^{(1)}(r_i - r_{01}^{(1)})^2}}{r_i} + B_{\text{HH}}^{(1)} \sum_{i=4}^6 \frac{e^{-\beta_{\text{HH}}^{(1)}(r_i - r_{01}^{(1)})^2}}{r_i} \quad (16)$$

with $0 \leq i, j, k \leq 2$, $i + j + k \leq 4$, $l + m + n \leq 2$, $i + j + k + l + m + n \geq 1$, and $1 \leq p \leq 2$. In Eq. (14), θ_0 is the ground-state equilibrium HNH bond angle which is fixed to the experimental value¹⁹ of 106.68 degrees; and $r_{01}^{(1)}$ is fixed to -0.2 \AA . There are a total of 64 linear coefficients, which include 62 $C_{ijklmnp}^{(1)}$ plus $B_{\text{NH1}}^{(1)}$ and $B_{\text{HH}}^{(1)}$, and five additional nonlinear parameters, $\gamma^{(1)}$, $r_0^{(1)}$, $\delta^{(1)}$, $\beta_{\text{NH1}}^{(1)}$, and $\beta_{\text{HH}}^{(1)}$, determined by fitting to the MC-QDPT energies.

First the values of the 69 fitting parameters specified above were obtained by fitting (discussed in section 4.3) to the data sets described in Section 2 with $E_{\text{D}} = 0$. Then E_{D} was included in Eq. (11) to constrain the analytical function to give the experimental dissociation limit and excitation energies and to eliminate unphysical values at very short N–H bond distances. The correction function used is:

$$E_{\text{D}}(r_1, r_2, r_3, r_4, r_5, r_6) = \Delta E_1^{(1)} + E_{\text{R}}(r_1, r_2, r_3) + \Delta E_2^{(1)} S_c(r_1, r_2, r_3), \quad (17)$$

where

$$E_{\text{R}}(r_1, r_2, r_3) = B_{\text{NH2}}^{(1)} \sum_{i=1}^3 \frac{e^{-\beta_{\text{NH2}}^{(1)}(r_i - r_{01}^{(1)})^2}}{r_i} \quad (18)$$

$$S_c(r_1, r_2, r_3) = \sum_{i=1}^3 \frac{1 + \tanh\left[\eta_1^{(1)}(r_i - 2r_e^{(1)})\right]}{2} \quad (19)$$

E_{R} is a function that adds a steep repulsive wall at very short N–H bond distances. The parameters $B_{\text{NH2}}^{(1)}$ and $\beta_{\text{NH2}}^{(1)}$ in the function E_{R} are manually adjusted so that there are no unphysical regions (with negative energies) at very short N–H bond distances. The parameter $r_e^{(1)}$ in the function S_c is fixed to the experimental N–H bond distance of 1.024 \AA of NH_2 ($\tilde{X}^2\text{B}_1$),²² while $\eta_1^{(1)}$ is manually adjusted so that the correction is largest when one N–H bond is broken but is negligible when the NH_3 geometry is close to the minimum equilibrium geometry. $\Delta E_1^{(1)}$ and $\Delta E_2^{(1)}$ are manually adjusted to reproduce the D_e of the NH_3 ($\tilde{X}^1\text{A}_1$) \rightarrow

$\text{NH}_2 (\tilde{\text{A}}^2\text{A}_1) + \text{H}$ reaction and to force the adiabatic V_1 energy of the minimum equilibrium geometry of NH_3 on the V_1 potential energy surface to be as close to zero as possible.

In the previous work, the U_{22} surface was fitted using a multiconfigurational representation involving two surfaces W_{11} and W_{22} .¹ In the present work, it is represented with just one configuration. The function form used is similar to that of U_{11} . However, to fit the excited-state minimum and the saddle point for the N–H bond dissociation on the U_{22} surface simultaneously, i, j and k are allowed to be equal to 3, but when they are equal to 3, function $B_i(r)$ is changed to

$$B_i(r_1, r_2, r_3) = e^{-\alpha^{(2)}(r_1 - r_{02}^{(2)})^2} e^{-\delta^{(2)}(r_2^2 + r_3^2)}, \quad i = 3. \quad (20)$$

When i, j , and k are less than or equal to 2, Eq. (13) is still used. Function $g(\theta_1, r_2, r_3)$ is changed to

$$g(\theta_1, r_2, r_3) = 1 + [\cos(\theta_0) - \cos(\theta_1)]^l e^{-\delta^{(2)}(r_2^2 + r_3^2)}. \quad (21)$$

The third difference between the functions used for U_{11} and U_{22} is that $h(p, \beta, r_1, r_2, r_3)$ is changed to

$$h(p, \beta, r_1, r_2, r_3) = 1 + \left\{ 1 - \cos \left[p \left(\beta - \frac{\pi}{2} \right) + \frac{1}{2} \frac{k^{(2)} \left(\beta - \frac{\pi}{2} \right)^2}{C^{(2)} + \left(\beta - \frac{\pi}{2} \right)^2} \right] \right\} e^{-\delta^{(2)}(r_1^2 + r_2^2 + r_3^2)}. \quad (22)$$

The reason for using Eq. (22) instead of Eq. (15) for U_{22} is that the harmonic vibrational frequency of the umbrella vibration motion for the excited-state minimum is too low at the level of theory used to prepare the data sets (see Table 4). This additional term effectively makes the surface steeper when the β angle deviates from the equilibrium value ($\pi/2$). The ranges of the i, j, k, l, m, n , and p values are slightly different from those used for U_{11} . The ranges for U_{22} are $0 \leq i, j, k \leq 3$, $0 \leq l, m, n \leq 2$, and $0 \leq p \leq 2$ with additional constraints that $i = j = k = l = m = n = 0$ when $p = 0$; $0 \leq l + m + n \leq 2$, $i + j + k + l + m + n \geq 1$, when $1 \leq p \leq 2$ and $0 \leq i + j + k \leq 3$; i, j , and $k \neq 2$, $0 \leq l + m + n \leq 2$, $i + j + k + l + m + n \geq 1$, when $1 \leq p \leq 2$ and $4 \leq i + j + k \leq 5$. For the U_{22} surface, θ_0 is fixed to the equilibrium value of 120 degrees, and $r_{01}^{(2)}$ is fixed to -0.2 \AA .

$B_{\text{NH1}}^{(2)}$, $\beta_{\text{NH1}}^{(2)}$, $B_{\text{HH}}^{(2)}$ and $\beta_{\text{HH}}^{(2)}$ are the same as those used for U_{11} , and $C^{(2)}$ is fixed to a small value of 0.05 rad². A total of 71 linear $C_{ijklmnp}^{(2)}$ coefficients and 6 nonlinear parameters, $\gamma^{(2)}$, $r_0^{(2)}$, $\delta^{(2)}$, $r_{02}^{(2)}$, $\alpha^{(2)}$, and $k^{(2)}$, are determined by fitting to the MC-QDPT energies ($k^{(2)}$ is a linear coefficient, but is treated as a nonlinear parameter in the fitting).

For the functions (Eqs. (17)–(19)) used to correct to the experimental dissociation limits after fitting to the MC-QDPT energies, the parameters $B_{\text{NH2}}^{(2)}$ and $\beta_{\text{NH2}}^{(2)}$ are manually adjusted. The parameters used in the function S_c are the same as those used for U_{11} , i.e., $\eta_1^{(2)} = \eta_1^{(1)}$ and $r_e^{(2)} = r_e^{(1)}$. $\Delta E_1^{(2)}$ and $\Delta E_2^{(2)}$ are manually adjusted to reproduce three experimental energy differences: 1) D_e for the $\text{NH}_3 (\tilde{X}^1A_1) \rightarrow \text{NH}_2 (\tilde{X}^2B_1) + \text{H}$ reaction; 2) T_e for the $\text{NH}_3 (\tilde{X}^1A_1) \rightarrow \text{NH}_3 (\tilde{A}^1A''_2)$ excitation; and 3) T_e for the $\text{NH}_2 (\tilde{X}^2B_1) \rightarrow \text{NH}_2 (\tilde{A}^2A_1)$ excitation.

For fitting the diabatic coupling surface U_{12} , the previous functional form¹ is used:

$$U_{12}(\mathbf{Q}) = \sum_{\substack{i \leq j \leq k \\ l \leq m \leq n \\ p}} C_{ijklmnp}^{(3)} \sum_{\alpha} B_{ijklmnp}^{(3)} (\mathbf{P}_{\alpha} \mathbf{Q}), \quad (23)$$

where

$$B_{ijklmnp}^{(3)}(\mathbf{Q}) = r_1^i r_2^j r_3^k F(r_1) F(r_2) F(r_3) \theta_1^l \theta_2^m \theta_3^n G(p, \beta), \quad (24)$$

with

$$F(r) = 1 - \tanh\left(r - r_0^{(3)}\right), \quad (25)$$

$$G(p, \beta) = \sin\left[p\left(\beta - \frac{\pi}{2}\right)\right], \quad (26)$$

where $0 \leq i, j, k \leq 2$, $0 \leq l, m, n \leq 2$, and $3 \leq p \leq 5$ with additional constraints $i + j + k \leq 2$, $l + m + n \leq 2$, and $i + j + k + l + m + n \geq 1$. There are a total of 45 linear $C_{ijklmnp}^{(3)}$ coefficients and only one nonlinear parameter, $r_0^{(3)}$, to be determined.

4.3. Fitting procedure

The linear coefficients were obtained by a linear least squares fitting routine for given nonlinear parameters. The nonlinear parameters are fitted using a simplex algorithm. To find the best parameters, random initial guesses were used, and the fitting procedure was repeated until

the error could not be lowered. The error minimized in the nonlinear fitting procedure is calculated by:

$$Err = \frac{\sum_{i=1}^{8227} w_i |E_i^F - E_i^{MC}|}{\sum_{i=1}^{8227} w_i} + \frac{\sum_{j=1}^{8227} \sum_{i=1}^{8227} w_i w_j (E_i^F - E_j^F) - (E_i^{MC} - E_j^{MC})}{\sum_{j=1}^{8227} \sum_{i=1}^{8227} w_i w_j}, \quad (27)$$

where w_i is the weight for the given energy, E_i^F is the energy given by the fitting function with the linear coefficients determined in the linear squares fitting with the same set of weights for the given nonlinear parameters, and E_i^{MC} is the energy to be fitted. For the fitting of the U_{11} surface, the weights are calculated by

$$w_i = \begin{cases} \left(\frac{3.1}{U_{11}(i)+0.1}\right)^{1.3}, & \text{when } \frac{3.1}{U_{11}(i)+0.1} < 1 \text{ and } r_1 < 2.0 \text{ \AA} \\ \left(\frac{3.1}{U_{11}(i)+0.1}\right)^{1.2}, & \text{when } \frac{3.1}{U_{11}(i)+0.1} \geq 1 \text{ and } r_1 < 2.0 \text{ \AA} \\ 3\left(\frac{3.1}{U_{11}(i)+0.1}\right)^{1.2}, & \text{when } U_{11}(i) \leq 7.0 \text{ eV, and } 2.0 \leq r_1 \leq 5.0 \text{ \AA}, \\ 35\left(\frac{3.1}{U_{11}(i)+0.1}\right)^{1.2}, & \text{when } U_{11}(i) \leq 7.0 \text{ eV, and } r_1 > 5.0 \text{ \AA} \\ \left(\frac{3.1}{U_{11}(i)+0.1}\right)^{1.3}, & \text{when } U_{11}(i) > 7.0 \text{ eV, and } r_1 \geq 2.0 \text{ \AA} \end{cases}, \quad (28)$$

where U_{11} is the MC-QDPT value in eV. Notice that extra weight was given to the dissociated points (that is, those with one N–H distance greater than 5.0 Å); this was done to force the function to have the right dissociation limit.

For the U_{22} surface, the weights used are

$$w_i = \begin{cases} \left(\frac{7.0}{U_{22}(i)}\right)^2 + \chi_i + \lambda_i, & \text{when } \frac{7.0}{U_{22}(i)} \geq 1 \text{ and } r_1 < 1.9 \text{ \AA} \\ 7\left(\frac{7.0}{U_{22}(i)}\right)^2, & \text{when } \frac{7.0}{U_{22}(i)} \geq 1.1 \text{ and } 1.9 \leq r_1 < 3.0 \text{ \AA} \\ 7.5\left(\frac{7.0}{U_{22}(i)}\right)^2, & \text{when } \frac{7.0}{U_{22}(i)} \geq 1.1 \text{ and } 3 \leq r_1 \leq 5.0 \text{ \AA} \\ 5\left(\frac{7.0}{U_{22}(i)}\right)^2, & \text{when } \frac{7.0}{U_{22}(i)} \geq 1 \text{ and } r_1 > 5.0 \text{ \AA} \end{cases}, \quad (29a)$$

otherwise

$$w_i = \begin{cases} \left(\frac{7.0}{U_{22}(i)}\right)^{1.6}, & \text{when } \frac{7.0}{U_{22}(i)} < 0.5 \\ \left(\frac{7.0}{U_{22}(i)}\right)^{1.3}, & \text{when } 0.5 \leq \frac{7.0}{U_{22}(i)} < 1, \\ \left(\frac{7.0}{U_{22}(i)}\right)^2, & \text{when } \frac{7.0}{U_{22}(i)} \geq 1 \end{cases} \quad (29b)$$

where the energies are in eV and χ_i and λ_i are functions of the three N–H distances chosen to add extra weight to the data points around the planar D_{3h} minimum and to the data points around the N–H dissociation transition state on the U_{22} surface:

$$\chi_i = 10e^{-50 \min[(r_i - r_t), i=1,3]}, \quad (30)$$

$$\lambda_i = 25e^{-50 \left(\frac{r_1 + r_2 + r_3}{3} - r_0\right)^2}, \quad (31)$$

where $r_t = 1.30 \text{ \AA}$ is the dissociating N–H distance in the transition state, and $r_0 = 1.05 \text{ \AA}$ is the equilibrium N–H distance of the planar D_{3h} minimum optimized by Dixon.³⁷

To fit the U_{12} surface, a uniform weighting scheme was used. It should be noted that since the functional form used for the U_{12} surface is an odd function of $(\beta - \pi/2)$, only those geometries with $\beta < \pi/2$ (7036 out of 8227) in the whole dataset were used to fit the U_{12} surface.

All the values for the linear coefficients and fitted nonlinear parameters are provided in the supporting information.³⁸

4.4. Availability

The fitted surfaces and coupling, including analytic gradients in Cartesian coordinates, have been deposited in the POTLIB³⁹ potential-energy surface library, which is freely available online.⁴⁰

5. Discussion

The discussion is organized as follows: In the next two paragraphs we discuss the accuracy of the preliminary fits before correcting the four critical energy differences (i.e., the

mean unsigned errors for the fits with $E_D = 0$). Then, we compare the properties of some stationary points obtained with the full functional forms (with $E_D \neq 0$) to the best available results, and finally we present mappings of the conical intersection seams.

The mean unsigned errors for preliminary fits are listed in Tables 6 and 7. Experimentally, the dissociated NH_2 ($\tilde{\text{A}}^2\text{A}_1$) + H is at least 6.902 eV above the bottom of the potential energy well of the NH_3 minimum on the adiabatic V_1 surface (the zero of energy for the current fitting). For fitting U_{11} , the MUE for energies between 0 and 7 eV is 0.11 eV, which we consider satisfactory. For fitting U_{22} , since we add extra weight to the points around the D_{3h} minimum and the dissociation transition state, the MUE of the fitted U_{22} energies between 5.7 and 6.7 eV is 0.05 eV. For all the energies below 7 eV, the MUE of the fitted U_{22} energies is 0.06 eV, which can also be considered satisfactory. For the fitting of the U_{12} surface, the MUE of the fitted U_{12} energies below 2 eV is 0.12 eV. The error is slightly larger than those of the U_{11} and U_{22} surfaces.

Having obtained the diabatic potential-energy surfaces and their coupling, the adiabatic surfaces V_1 and V_2 can be obtained by¹

$$V_{1(2)} = \frac{1}{2} \left\{ U_{11}(\mathbf{Q}) + U_{22}(\mathbf{Q}) \mp \sqrt{[U_{11}(\mathbf{Q}) - U_{22}(\mathbf{Q})]^2 + 4U_{12}(\mathbf{Q})^2} \right\}. \quad (32)$$

Although our fit does not use V_1 and V_2 energies at all, the MUE errors of the fitted V_1 and V_2 energies are reasonably good (Table 8): The MUE of the fitted V_1 energies below 7 eV is 0.10 eV, and that of the fitted V_2 energies below 7 eV is just 0.05 eV.

To characterize the quality of the fitted potential energy surfaces, we have compared the geometries and vibrational frequencies of some important stationary points to the best previous results. The results are presented in Tables 4 and 5. The N–H bond distance of NH_3 minimum on the adiabatic ground state is 1.016 Å, very close to the experimental value of 1.012 Å.¹⁹ The HNH bond angle is just half degree smaller than the experiment value. Although we did not explicitly fit to the experimental harmonic vibrational frequencies of the ground state NH_3 minimum, the fitted frequencies agree reasonably well with experiment. Table 8 indicates that

the fitting around the ground-state minimum is very accurate, with an MUE of just 0.01 eV for the energy range 0–1 eV. Therefore, the discrepancy between the vibrational frequencies given by the fitted function and those experimental values must be due mainly to the inaccuracy of the MC-QDPT method. The N–H bond distance of the NH_3 minimum on the adiabatic excited state $\tilde{\text{A}}^1\text{A}''_2$ is 1.039 Å, in good agreement with the experimental value of 1.055 Å.²⁴ The harmonic vibrational frequencies are also in fairly good agreement with experimental values.

We optimized the transition state for N–H dissociation on the diabatic U_{22} surface using the optimizer in the *Gaussian 03* program¹⁷ with its “external” functionality. The dissociating N–H bond distance at the optimized saddle point with the current analytical function is slightly shorter than the best available value of 1.298 Å²⁵ by 0.009 Å, while the other two N–H bond distances are very close to the literature values.²⁵ For the dynamics simulation of the photodissociation of ammonia, a property that is probably more important is the barrier height. The classical barrier height relative to energy of the planar NH_3 ($\tilde{\text{A}}^1\text{A}''_2$) minimum given by the analytical function is 0.22 eV, 0.07 eV lower than the best literature value of 0.29 eV.²⁵

For the minima of the ground-state and the first singlet excited-state NH_2 , the agreement between the current fitting and experiments is very good. The agreement between the fitted vibrational frequencies and the best available values is also fairly good. It should be noted that since we manually adjust the function to give the four experimental energy differences, they are exactly the same as the experimental values. The U_{11} (V_2) and U_{22} (V_1) energies of the NH_2 are degenerate at linear geometries. Since U_{11} and U_{22} potential surfaces were fitted with different functional forms, it is hard for the analytical function to produce identical U_{11} and U_{22} energies for the linear geometries. In Figure 3 the U_{11} and U_{22} energies of NH_2 were plotted respect to the HNH angle. The current analytical functions give almost the same U_{11} and U_{22} energies when NH_2 is linear.

As pointed out in the previous paper on the NH_3 PESs,¹ the conical intersection is at most four dimensional, since when V_1 equals to V_2 , U_{22} must equal to U_{11} , and U_{12} must vanish according to Eq. (32). Apparently, U_{12} vanishes only for planar structures. In Figure 4, we

present one-dimensional plots of U_{11} and U_{22} as functions of one N–H distance (r_1) for the planar structures with the constraint of D_{3h} (Fig. 4a), C_{2v} (Fig. 4b), and C_s (Fig. 4c) symmetry. The geometrical parameters other than r_1 are optimized to minimize U_{22} subject to the specific symmetry. The geometries and energies of the three conical intersection points in Figure 4 are listed in Table 9. The plot in Fig. 4a shows two conical intersection points. The one with higher energy corresponds to the dissociated N + 3H, which is qualitatively correct. The geometry of the one presented in Table 9 is the one with lower energy. One can see that the conical intersection with D_{3h} symmetry has the shortest N–H bond distance, while that with C_{2v} symmetry has the longest N–H bond distance. These scans, though, do not locate the lowest-energy conical intersections for these symmetries. To find the lowest-energy conical intersection we minimize the function

$$F(\mathbf{Q}) = \frac{1}{2}[U_{11}(\mathbf{Q}) + U_{22}(\mathbf{Q})] + \mu[U_{11}(\mathbf{Q}) - U_{22}(\mathbf{Q})]^2, \quad (33)$$

where μ is an adjustable parameter. As μ is increased, the geometry that minimizes this function (for planar geometries) is the lowest-energy conical intersection. We varied μ from 1000 E_h^{-1} to 50000 E_h^{-1} until the final geometry converges. The final resulting geometry is given in Table 9.

Yarkony found several conical intersection geometries with both C_{2v} and C_s symmetries.⁴¹ The lowest-energy conical intersection has C_{2v} symmetry. The one we obtained is higher in energy by 0.06 eV. This is because the present fitting exactly reproduces the dissociation energies and the $\text{NH}_3(\tilde{X}^1A_1) \rightarrow \text{NH}_3(\tilde{A}^1A''_2)$ equilibrium excitation energy (T_e), which are all higher than the corresponding theoretical values.⁴¹ Therefore, the whole U_{22} surface, and the U_{11} surface with large N–H bond distance, are all shifted upwards, and the energies of the conical intersections are shifted upwards accordingly.

In Figure 5 we plot the three-dimensional diabatic and adiabatic surfaces varying one N–H distance (r_1) and the trisector angle β while keeping the two other N–H bond distances fixed at the planar equilibrium geometry (1.039 Å) on the U_{22} surface and the three projected angles ($\theta'_1, \theta'_2, \theta'_3$) on the plane fixed at 120 degrees. Fig. 5a shows the crossing between the U_{11} and

U_{22} surfaces. Although U_{11} crosses U_{22} for both planar and nonplanar geometries, only the former crossings are conical intersections since U_{12} is not zero for nonplanar structures.

In Figure 6 we present three-dimensional (Fig. 6a) and contour (Fig. 6b) plots of the conical intersection seam with the constraint of C_{2v} symmetry. The scan is carried out with the external function of the *Gaussian* 03 program with other geometrical parameters optimized on the U_{22} surface. Actually, only one parameter, i.e., one H-N-H angle, is allowed to relax under the constraint of C_{2v} symmetry. Fig. 6a shows the seam of the crossing between the U_{11} and U_{22} surfaces. Fig. 6b is a contour plot of the differences between the two diabatic energies ($U_{22} - U_{11}$). At shorter r_1 , U_{22} is higher in energy while at longer r_1 , U_{22} becomes the ground electronic state. Since U_{12} vanishes for the planar structures, and the geometries are all planar, the crossing seam between the U_{11} and U_{22} surfaces is the location of the conical intersections. The plots indicate that the r_1 distance of the conical intersection geometries becomes shorter if the other two N-H distances become longer.

The photodissociation patterns of the excited NH_3 ($\tilde{A}^1A''_2$) strongly depend on the vibrational mode being excited.⁴² Figure 7 shows one-dimensional cuts through the potential surfaces for slightly nonplanar geometries with a trisector angle of 77.4 degrees in both C_s and C_1 symmetries; U_{22} is dashed. Fig. 7b and Fig. 7c correspond to symmetrical stretching of the N-H bonds while Fig. 7d corresponds to an asymmetrical stretching of the N-H bonds.

6. Conclusions

In this paper we have improved the application of the fourfold way direct diabaticization scheme to ammonia. The improvements in the diabaticization are:

- 1) We define a geometry-dependent p orbital as a globally appropriate reference diabatic molecular orbital.
- 2) We use carefully selected dominant configuration lists that include the dominant character of the diabatic configuration state functions at a global set of geometries.

Using the diabatic energies and couplings calculated with the new diabaticization scheme, we have refitted the first two diabatic potential-energy surfaces, U_{11} and U_{22} , and their coupling, U_{12} , with improved functional forms and an improved fitting scheme. The improvements of the functional forms are:

- 1) A correction function is used to reproduce the four most important experimental energy differences exactly: a) the dissociation energy D_e for the dissociation of ammonia to the ground state of products; b) the dissociation energy D_e for the dissociation of ammonia to the excited state of products; c) the excitation energy T_e for $\text{NH}_3 (\tilde{X}^1\text{A}_1) \rightarrow \text{NH}_3 (\tilde{\text{A}}^1\text{A}_2^-)$; and d) the excitation energy T_e for $\text{NH}_2 (\tilde{X}^2\text{B}_1) \rightarrow \text{NH}_2 (\tilde{\text{A}}^2\text{A}_1)$.
- 2) We use a new function $g(\theta_i, r_j, r_k)$ for fitting the dependence on bond angle because it has better behavior for $\theta_i = \pi$.
- 3) The U_{22} potential-energy surface is fitted with just one functional form, as for U_{11} , and the number of terms in the polynomials used in the fitting functions for both U_{11} and U_{22} are greatly reduced. Therefore the new analytical potential can be evaluated significantly faster than the previous function.

7. Acknowledgments

The authors are grateful to Shikha Nangia and Ahren Jasper for assistance and to Mark Gordon for permission to port the ISA algorithm from GAMESS to HONDOPLUS. This work was supported in part by the National Science Foundation through grant no. CHE03-49122.

References:

- 1 Nangia S, Truhlar DG (2006) *J Chem Phys* 124:124309
- 2 Nakamura H, Truhlar DG (2001) *J Chem Phys* 115:10353
- 3 Nakamura H, Truhlar DG (2002) *J Chem Phys* 117:5576
- 4 Nakamura H, Truhlar DG (2003) *J Chem Phys* 118:6816
- 5 Atchity GJ, Ruedenberg K (1993) *J Chem Phys* 99:3790
- 6 Ruedenberg K, Atchity GJ (1993) *J Chem Phys* 99:3799
- 7 Nakano H (1993) *J Chem Phys* 99:7983
- 8 Nakano H (1993) *Chem Phys Lett* 207:372
- 9 Nakano H, Nakajima T, Tsuneda T, Hirao K (2001) *J Mol Struct* 573:91
- 10 Ruedenberg K, Sundberg R (1976) In Calais, J-L, Goscinski O, Linderberg J, Öhrn, Y (Eds) *Quantum Science*. Plenum, New York, p 505.
- 11 Roos, BO (1987) *Adv Chem Phys* 69:399
- 12 Witek HA, Choe Y-K, Finley JP, Hirao K, (2002) *J Comput Chem* 10:957
- 13 Krishnan R, Binkley JS, Seeger R, Pople JA (1980) *J Chem Phys* 72:650
- 14 Frisch MJ, Pople JA, Binkley JS (1984) *J. Chem Phys* 80:3265
- 15 Nakamura H, Xidos JD, Thompson JD, Li J, Hawkins GD, Zhu T, Lynch BJ, Volobuev Y, Rinaldi D, Liotard DA, Valero R, Cramer CJ, Truhlar DG, HONDOPLUS v4.7, based on HONDO-v99.6; University of Minnesota, Minneapolis (2006).
- 16 Schmidt MW, Baldridge KK, Boatz JA, Elbert ST, Gordon MS, Jensen JH, Koseki S, Matsunaga N, Nguyen KA, Su SJ, Windus TL, Dupuis M, Montgomery JA (1993) *J Comput Chem* 14:1347
- 17 Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Montgomery Jr. JA, Vreven T, Kudin KN, Burant JC, Millam JM, Iyengar SS, Tomasi J, Barone V,

- Mennucci B, Cossi M, Scalmani G, Rega N, Petersson GA, Nakatsuji H, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Klene M, Li X, Knox JE, Hratchian HP, Cross JB, Bakken V, Adamo C, Jaramillo J, Gomperts R, Stratmann RE, Yazyev O, Austin AJ, Cammi R, Pomelli C, Ochterski JW, Ayala PY, Morokuma K, Voth GA, Salvador P, Dannenberg JJ, Zakrzewski VG, Dapprich S, Daniels AD, Strain MC, Farkas O, Malick DK, Rabuck AD, Raghavachari K, Foresman JB, Ortiz JV, Cui Q, Baboul AG, Clifford S, Cioslowski J, Stefanov BB, Liu G, Liashenko A, Piskorz P, Komaromi I, Martin RL, Fox DJ, Keith T, Al-Laham MA, Peng CY, Nanayakkara A, Challacombe M, Gill PMW, Johnson B, Chen W, Wong MW, Gonzalez C, and Pople JA, *Gaussian 03*, Revision D.01, Gaussian, Inc., Wallingford CT, 2004.
- 18 Léonard C, Handy NC, Carter S, Bowman JM (2002) *Spectrochim Acta Part A* 58:825
- 19 Landolt-Börnstein, *Numerical Data and Function Relationships in Science and Technology* II(7,2) (Springer, New York, 1976). Cited in Pesonen J, Miani A, Halonen L (2001) *J Chem Phys* 115:1243
- 20 Coy SL, Lehmann KK (1989) *Spectrochim Acta Part A* 45:47
- 21 Martin JML, Lee TJ, Taylor PR (1992) *J Chem Phys* 97:8361
- 22 Handy NC, Carter S, Colwell SM (1999) *Mol Phys* 96:477
- 23 Mordaunt DH, Dixon RN, Ashfold MNR (1996) *J Chem Phys* 104:6472
- 24 Henck SA, Mason MA, Yan WB, Lehmann KK, Coy SL (1995) *J Chem Phys* 102:4772
- 25 Bach A, Hutchison JM, Holiday RJ, Crim FF (2002) *J Chem Phys* 116:9315
- 26 Morino I, Kawaguchi KJ (1997) *Mol Spectrosc* 182:428
- 27 Gabriel W, Chambaud G, Rosmus P, Carter S, Handy NC (1994) *Mol Phys* 81:1445
- 28 Jacox ME (1994) *J Phys Chem Ref Data Monograph* 3

- 29 Herzberg G, *Molecular Spectra and Molecular structure*, Vol. 3 (Van Nostrand Reinhold, New York, 1966). Cited in McCarthy MI, Rosmus P, Werner H-J, Botschwina P, Vaida V (1987) *J Chem Phys* 86:6693
- 30 As reported in <http://webbook.nist.gov/chemistry>
- 31 Ruscic B, Boggs JE, Burcat A, Császár, AG, Demaison J, Janoschek R, Martin JML, Morton ML, Rossi MJ, Stanton JF, Szalay PG, Westmoreland PR, Zabel F, Bérces T (2005) *J Phys Chem Ref Data* 34:573 and references therein
- 32 Hargiss LO, Ermler WC (1988) *J Phys Chem* 92:300
- 33 Gatti F, Lung C, Leforestier C, Chapuisat X (1999) *J Chem Phys* 111:7236
- 34 Demaison J, Margulès L, Boggs JE (2003) *Phys Chem Chem Phys* 5:3359
- 35 Barone V (2004) *J Chem Phys* 120:3059
- 36 Douglas AE (1963) *Discuss Faraday Chem Soc* 35:158
- 37 Dixon RA (1996) *Mol Phys* 88:949
- 38 Supporting information is available via the Theo Chem Acc journal homepage
- 39 Duchovic RJ, Volobuev Y L, Lynch GC, Truhlar DG, Allison TC, Wagner F, Garrett BC, Corchado JC (2002) *Comput Phys Commun* **144**:169; (2004) **156**:319(E).
- 40 Duchovic RJ, Volobuev Y L, Jasper AW, Lynch GC, Truhlar DG, Allison TC, Wagner F, Garrett BC, Espinosa-García J, Corchado JC POTLIB-online, <http://comp.chem.umn.edu/potlib>
- 41 Yarkony DR (2004) *J Chem Phys* 121:628
- 42 Bach A, Hutchison JM, Holiday RJ, Crim FF (2003) *J Phys Chem A* 107:10490

Table 1 Correlation of DMOs between the different NH₃ geometries by the threefold way.^a

β (deg) ^b	Small N–A distance (NH ₃)	Intermediate N–A distance (NH ₃)	Large N–A distance (NH ₂ + H)
$\beta = 90^c$	$\sigma_{\text{N–A}} p_z R$	$\sigma_{\text{N–A}} p_z R$	$sp^2 p_z h$
$75 < \beta < 90^d$	$\sigma_{\text{N–A}} n R$	$\sigma_{\text{N–A}} n R$	$sp^2 p_z h$
$\beta < 75^d$	$\sigma_{\text{N–A}} n R$	$n \sigma_{\text{N–A}} R$	$sp^2 p_z h$

^a Each entry in the table is an ordered list of three diabatic molecular orbitals; in comparing entries, the first orbital of one entry correlated with the first of another, the second with the second, and the third with the third. For small and intermediate N–A, the z direction is perpendicular to the NBC plane, sp^2 denotes an in-plane nonbonding orbital of NH₂, where “plane” refers to the NBC plane, h denotes an orbital localized on the dissociated hydrogen, and $\sigma_{\text{N–A}}$ is a bonding orbital.

^b β is the angle between an N–H bond axis and the trisector of all the N–H bond axes (see Fig. 2).

^c Planar

^d Nonplanar

Table 2 Correlation of DMOs between the different NH₃ geometries by the fourfold way ($\lambda = 1$).

	Small N–A distance (NH ₃)	Intermediate N–A distance (NH ₃)	Large N–A distance (NH ₂ + H)
Planar	$\sigma_{\text{N–A}} p_z R$	$\sigma_{\text{N–A}} p_z R$	$sp^2 p_z h$
Nonplanar	$\sigma_{\text{N–A}} n R$	$\sigma_{\text{N–A}} n R$	$sp^2 p_z h$

Table 3 Coordinate and diabatic energy ranges for all the computed points.

	r_2 (Å)	r_3 (Å)	r_1 (Å)	θ'_1 (deg)	θ'_2 (deg)	θ'_3 (deg)	β (deg)	U_{11} (eV)	U_{22} (eV)
Regular grid	0.8-1.5	0.8-1.5	0.8-5.0	90-150	60-150	60-150	30-90	0.1-30.4	4.7-37.3
GS min	1.0-1.1	1.0-1.1	1.0-1.1	87-153	87-153	87-153	60-90	0.1-2.3	5.9-8.2
ES N–A saddle point	0.99-1.05	0.99-1.05	1.05-1.95	102-153	105-130	75-155	75-90	0.4-5.6	5.4-8.9
ES min I	0.9-1.3	0.9-1.5	0.9-1.5	105-145	60-170	60-170	55-90	0.4-7.3	5.9-13.7
ES min II	1.03-1.07	1.00-1.07	1.0-1.07	117-123	116-123	116-123	86-92	0.3-0.6	5.8-6.0
NH ₂ (\tilde{X}^2B_1) + H	0.95-2.80	0.95-5.00	10.0	90	130-170	100-180	90	6.3-14.7	4.8-14.7
NH ₂ (\tilde{A}^2A_1) + H	0.95-2.80	0.95-5.00	10.0	90	90-130	100-180	90	6.1-14.7	5.4-14.7
Repulsive I	0.8-1.5	1.02-1.5	1.02-1.5	10-180	10-180	10-180	40-90	14-103	15-106
Repulsive II	0.5-0.6	0.5-0.7	0.6-1.5	10-150	30-180	40-150	10-90	26-134	32-138
Repulsive III	1.02	1.02	0.2-5.0	120	120	120	65	0-217	5-224
Repulsive IV	1.02	1.02	1.02	124-179	122-152	115-17	54-64	0.2-29.3	6.8-34.8
Repulsive V	0.2-0.8	0.4-0.8	0.2-0.4	10-40	10-40	10-40	10-20	302-305	309-312
Long N–A distance	1.04	1.04	3.3-4.1	60-180	60-150	60-150	30-90	6.2-9.2	4.9-6.4

Table 4 State-specific CASSCF(8,7)/6-311++G(3df, 3pd) geometries (Å and degree) and vibrational frequencies (cm⁻¹), state-averaged MC-QDPT energies (eV), experimental data, and analytical function results for NH₃. Zero-point energies corrected for anharmonicity are given in parentheses, following harmonic values. D_0 and T_0 are given in brackets.

Quantity	Present Fit			Theoretical			Best Estimate ^a				
V_1											
Equilibrium geometry (C_{3v})											
r_1	1.016			1.023			1.012 ¹⁹				
θ_1	106.1			105.0			106.7 ¹⁹				
Vibrational frequencies											
ω_1 (a_1)	ω_2 (a_1)	3397.4	1235.2	3358.7	1144.4	3484.6 ²⁰	1084.1 ^{b,21}				
ω_3 (e)	ω_4 (e)	3628.0	1750.2	3473.8	1700.2	3623.7 ²⁰	1677.8 ²⁰				
ZPE		0.954			0.921			0.941 (0.925 ²²)			
D_e for NH ₃ (\tilde{X}^1A_1) → NH ₂ (\tilde{X}^2B_1) + H		5.016			4.790			5.016 [4.602 ^{b,23}]			
V_2											
Equilibrium geometry (D_{3h})											
r_1	1.039			1.036			1.055(8) ²⁴				
Vibrational frequencies											
ω_1 (a_1')	ω_2 (a_2'')	2776.9	891.2	2959.3	314.6	2870	892				
ω_3 (e')	ω_4 (e')	3041.2	1329.9	3213.6	1437.1	3020	1110 ²⁴				
ZPE		0.769			0.780			0.745 ²⁴ (0.724 ²⁴)			
T_e for NH ₃ (\tilde{X}^1A_1) → NH ₃ (\tilde{A}^1A_2'')		5.923			5.870			5.923 [5.722 ²⁴]			
Saddle point geometry (C_{2v})											
r_1	r_2	r_3	1.289	1.040	1.040	1.274	1.038	1.038	1.298	1.040	1.040
θ_1	θ_2	θ_3	112.1	123.9	123.9	113.4	123.3	123.3	112.4	123.8	123.8 ^{b,25}
Vibrational frequencies											
ω_1 (a_1)	ω_2 (a_1)	ω_6 (a_1)	3056.9	1417.6	1866.7i	3115.4	1503.5	2608.8i			
ω_3 (b_1)	ω_4 (b_2)	ω_5 (b_2)	941.1	3078.2	569.9	916.6	3333.4	664.8			
Saddle point energy without ZPE		6.14			6.11			6.19 ^{b,25}			
Classical barrier height		0.22			0.24			0.29 ^{b,25}			

^a From experiments except where indicated otherwise. ^b Theoretical value.

Table 5 State-specific CASSCF(8,7)/6-311++G(3df, 3pd) geometries (Å and degree) and frequencies (cm⁻¹), state-averaged MC-QDPT energies (eV), experimental data, and analytical function results for NH₂. Zero-point energies corrected for anharmonicity are given in parentheses, following harmonic values.

Quantity			Present Fit			Theoretical			Best Estimate		
V₁											
NH ₂ (\tilde{X}^2B_1) equilibrium geometry											
R_1			1.034			1.037			1.024 ²⁶		
θ_1			103.1			101.8			103.4 ²⁶		
Vibrational frequencies											
ω_1 (a ₁)	ω_2 (a ₁)	ω_3 (b ₂)	3212.0	1600.7	3553.1	3244.5	1556.3	3340.5	3374.2	1523.5	3481.2 ²⁷
ν_1 (a ₁)	ν_2 (a ₁)	ν_3 (b ₂)			3219	1497	3301 ²⁸
ZPE			0.519			0.505			0.519 ²⁷ (0.511 ²⁷)		
V₂											
NH ₂ (\tilde{A}^2A_1) equilibrium geometry											
r_1			0.992			1.000			1.004 ²⁹		
θ_1			143.4			142.8			144 ²⁹		
Vibrational frequencies											
ω_1 (a ₁)	ω_2 (a ₁)	ω_3 (b ₂)	3629.1	1008.4	3889.1	3592.8	1054.1	3897.0	3635	964	3953 ²⁷
ν_1 (a ₁)	ν_2 (a ₁)	ν_3 (b ₂)			3486.3	1158.1	3780.0 ²⁷
ZPE			0.529			0.530			0.530 ²⁷ (0.529 ^a)		
NH ₂ (\tilde{X}^2B_1) → NH ₂ (\tilde{A}^2A_1) excitation energy											
T_0					1.379 ³⁰		
T_e			1.361			1.410			1.361		

^a Estimated using Eq. (8).

Table 6 Mean unsigned error of the fitted U_{11} and U_{22} energies with respect to the state-averaged MC-QDPT energies for several energy ranges (all energies in eV).

U_{11}			U_{22}		
ΔE	N^a	MUE	ΔE	N^a	MUE
0–1	806	0.01	0–5	198	0.03
0–2	1628	0.02	0–6	1088	0.04
0–3	1906	0.03	0–7	3771	0.06
0–4	2147	0.04	0–8	4930	0.11
0–5	2627	0.06	0–9	6202	0.17
0–6	3198	0.09	0–10	6914	0.21
0–7	4022	0.11	0–11	7304	0.23
0–8	4896	0.13	0–12	7551	0.24
0–9	5721	0.16	0–13	7708	0.25
0–10	6712	0.20	0–14	7806	0.26

^a Number of data points in the corresponding energy range, ΔE .

Table 7 Mean unsigned error of the fitted U_{12} energies to the state-averaged MC-QDPT energies for several energy ranges (all energies in eV).

ΔE	N^a	MUE
0–0.2	1992	0.08
0–0.4	3136	0.09
0–0.6	4034	0.09
0–0.8	4802	0.09
0–1.0	5518	0.10
0–1.2	5983	0.10
0–1.4	6334	0.11
0–1.6	6616	0.12
0–1.8	6802	0.12
0–2.0	6914	0.12

^a Number of data points in the corresponding energy range, ΔE .

Table 8 Mean unsigned error (eV) of the fitted V_1 and V_2 energies to the state-averaged MC-QDPT energies for several energy ranges (all energies in eV).

V_1			V_2		
ΔE	N^a	MUE	ΔE	N^a	MUE
0–1	829	0.01	0–5	0	-
0–2	1634	0.02	0–6	281	0.04
0–3	1948	0.03	0–7	2560	0.05
0–4	2258	0.05	0–8	3246	0.08
0–5	3170	0.07	0–9	4362	0.14
0–6	4127	0.09	0–10	5690	0.23
0–7	5503	0.10	0–11	6669	0.28
0–8	6367	0.12	0–12	7244	0.31
0–9	7338	0.14	0–13	7599	0.32
0–10	7680	0.15	0–14	7730	0.33

^a Number of data points in the corresponding energy range, ΔE .

Table 9 Geometries (Å and degrees) and energies (eV) of the conical intersections in Figure 4, and the lowest-energy conical intersections given by the current analytical functions and the literature.

	r_1	r_2	r_3	θ_1	θ_2	θ_3	V_1	V_2
Fig. 4a	1.628	1.628	1.628	120.0	120.0	120.0	8.74	8.74
Fig. 4b	1.952	1.028	1.028	104.0	128.0	128.0	5.17	5.17
Fig. 4c	1.832	1.002	1.050	102.2	164.0	93.8	5.27	5.27
C_{2v}^a	1.990	1.023	1.023	107.9	126.0	126.0	5.16	5.16
C_{2v}^b	1.955	1.021	1.021	110	125	125	5.11	5.11

^a The lowest-energy conical intersection geometry given by the current analytical functions.

^b From Ref. 41.

FIGURE CAPTIONS

Figure 1. Pictorial representation of the NH_3 system with p_x - and p_z -type DMOs which mix as the N–A distance and the trisector angle change. See text for a detailed explanation.

Figure 2. Internal coordinates used for fitting the diabatic surfaces and their couplings. r_1 , r_2 and r_3 are the N–A, N–B and N–C bond distances, respectively. r_4 , r_5 and r_6 are the H_B – H_C , H_A – H_C and H_A – H_B bond distances, respectively. θ_1 is the bond angle $\text{H}_\text{B}\text{NH}_\text{C}$, θ_2 is the bond angle $\text{H}_\text{A}\text{NH}_\text{C}$, θ_3 is the bond angle $\text{H}_\text{A}\text{NH}_\text{B}$, and β is the trisector angle. The prime denotes the projection of the atoms on a plane perpendicular to the NN' direction.

Figure 3. One-dimensional potential-energy profiles of the NH_2 with the two N–H bond distances fixed at 1.02 Å. The solid line is U_{11} , and the dashed line is U_{22} .

Figure 4. One-dimensional potential-energy profiles of the relaxed scans on the diabatic U_{22} PES of the planar NH_3 with the constraint of: (a) D_{3h} symmetry; (b) C_{2v} symmetry; (c) C_s symmetry. The solid line is U_{11} , and the dashed line is U_{22} .

Figure 5. Three-dimensional plots of (a) the U_{11} and U_{22} diabatic PESs showing the diabatic crossing of the two surfaces and (b) the V_1 and V_2 adiabatic potential-energy surfaces of ammonia showing the conical intersection along the N–A stretch (r_1) and the inversion angle β . The other two N–H bond distances are fixed at 1.039 Å, and the three projected bond angles ($\theta'_1, \theta'_2, \theta'_3$) are fixed at 120 degrees.

Figure 6. (a) Three-dimensional plot of the U_{11} and U_{22} diabatic PESs and (b) the contour plot of their differences ($U_{22} - U_{11}$) showing the diabatic crossing of the two surfaces of planar ammonia with the constraint of C_{2v} symmetry. Other geometrical parameters are allowed to relax on the U_{22} PES with the constraint of C_{2v} symmetry.

Figure 7. V_2 , U_{11} , U_{22} , and V_1 vs. r_1 with $\theta_1 = 120$ deg, $\theta_2 = \theta_3 = 113$ deg, (a) $r_2 = r_3 = 1.04$ Å, (b) $r_2 = r_3 = 0.90$ Å, (c) $r_2 = r_3 = 1.19$ Å, (d) $r_2 = 0.90$ Å, $r_3 = 1.19$ Å.

Figure 1




Figure 2




Figure 3







Figure 4



(a)




(b)




(c)

Figure 5



(a)



(b)

Figure 6








Figure 7




(a)



(b)



(c)



(d)
